

(12) **UK Patent Application** (19) **GB** (11) **2 336 371** (13) **A**

(43) Date of A Publication 20.10.1999

(21) Application No 9807664.9

(22) Date of Filing 14.04.1998

(71) Applicant(s)  
**Reckitt & Colman Inc**  
(Incorporated in USA - Delaware)  
1655 Valley Road, Wayne, New Jersey 07474,  
United States of America

(72) Inventor(s)  
**Robert Zhong Lu**  
**Ralph Edward Rypkema**  
**Andrew Arno Kloeppel**  
**Dennis Thomas Smialowicz**  
**Nick Nanavati**  
**Mike Love**  
**Robert William Bogart**  
**Frederic Albert Taraschi**  
**Karen Ann McCue**  
**Diane Joyce Burt**

(51) INT CL<sup>6</sup>  
**C11D 3/00**

(52) UK CL (Edition Q )  
**C5D DHX D120 D121 D123 D124 D162 D165 D166 D182**

(56) Documents Cited  
**GB 2309706 A WO 95/31962 A1 US 4755327 A**

(58) Field of Search  
**UK CL (Edition Q ) C5D DHX DHZ DJX D162**  
**INT CL<sup>6</sup> C11D 3/00**  
**Online: WPI, CLAIMS**

(74) Agent and/or Address for Service  
**Martin N Dale**  
**Reckitt & Colman Plc, Group Patents Department,**  
**Dansom Lane, HULL, HU8 7DS, United Kingdom**

(54) Abstract Title  
**Aqueous disinfecting and cleaning composition**

(57) Aqueous disinfecting and cleaning compositions and concentrates which are efficacious against gram positive and gram negative bacteria, have relatively low volatile organic content ("VOC") and are surprisingly mild to the user of the compositions. The compositions include a quaternary ammonium compound as its primary germicidal active agent, have a low content of active constituents, and do not include organic solvents such as alcohols, glycols, or glycol ethers in significant amounts.

**GB 2 336 371 A**

## IMPROVEMENTS IN OR RELATING TO ORGANIC COMPOSITIONS

5

The present invention relates to improvements in cleaning compositions. More particularly the present invention is directed to improved cleaning compositions which find particular use in hard surface cleaning and disinfecting applications.

Certain hard surface formulations are known. For example, in US 3539520 to Cantor et al. are illustrated certain 'detergent-sanitizer' compositions. There are essentially based on aqueous mixtures of quaternary ammonium compounds as germicidal active agents in conjunction with nonionic surfactants based on alkoxy block copolymers, and in particular, compounds based on ethoxy/propoxy block copolymers. Therein Cantor notes that such nonionic surfactants surprisingly do not interfere with the germicidal effect of the quaternary ammonium compounds, and copious examples illustrate the utility of these specific nonionic surfactants as opposed to other classes of nonionic surfactants. Cantor notes, but however does not illustrate, any significant cleaning testing or results in his compositions, and in fact teaches away from the use of these particular nonionic surfactant compounds based on ethoxy/propoxy block copolymers in conjunction with other classes of nonionic surfactants. Cantor is also wholly silent as to the dermal and ocular irritation characteristics of his compositions.

More recently, in US 5454984 to Graubart et al. are recited all-purpose aqueous cleaning compositions which also include quaternary ammonium compounds as germicidal active agents, in conjunction with non-ionic surfactants which are desirably a ternary non-ionic surfactant system which includes three different nonionic surfactants. None of these nonionic surfactants of the ternary system are based on ethoxy/propoxy block copolymers. Further, the recited aqueous all-purpose aqueous cleaning compositions include an appreciable amount of an organic solvent constituent, believed to significantly facilitate the soil loosening and overall cleaning effects of these cleaning compositions.

Certain patents have recognized the fact that aqueous compositions containing quaternary ammonium compounds which provide a sanitizing benefit advantageously include one or more chemical compounds which function to mitigate such compositions, particularly with respect to the potential for ocular irritation. For example, in US Patent 4336151 to Like et al. therein are disclosed certain materials which are useful as irritation mitigants including certain ethoxylated cocodiethanolamides, certain polyoxyethylenes, certain hydrolyzed animal proteins, allantoin, 1,6-hexylene glycol, stearyl dimethylamine oxide, certain dextrose sugars and imidazole. US Patent 5547990 to Hall et al. discusses further irritation mitigants based on certain substituted imidazoline amphoteric surfactants as being useful in conjunction with didecyl dimethyl ammonium chloride, although this effect was not found with other types of quaternary ammonium compounds.

The contents of the US Patent documents indicated above are incorporated herein by reference.

Notwithstanding advantageous known art formulations, there yet remains a real and continuing need in the art for improved cleaning and disinfecting compositions in general, and in specific such compositions which provide at least one, but feature a plurality of the following characteristics: low volatile organic content, low irritancy to the end user of the composition, phase stability in storage (both at freeze-thaw, room temperature (i.e., 20°C) and elevated temperature (i.e., 40°C) conditions), ease of fabrication, low cost, efficacy against gram positive bacteria, efficacy against gram negative bacteria, good cleaning characteristics, and relatively low percentages actives required in such an aqueous formulation.

The compositions of the invention are aqueous disinfecting and cleaning compositions and concentrates thereof which are effective cleaning compositions and are efficacious as disinfecting compositions against gram positive and gram negative bacteria, have relatively low volatile organic content ("VOC") and are mild to the user of the compositions. That these results are concurrently achieved with a composition which includes a quaternary ammonium compound as its primary germicidal active agent is surprising, and indicates a synergistic effect not apparent from the prior art.

These compositions also provide good cleaning and disinfecting properties with low amounts of active constituents, and according to certain preferred embodiments do not include organic solvents such as low molecular weight alcohols, glycols or glycol ethers, in significant amounts, i.e., amounts in excess of about 1%wt and more.

5 In accordance with a first aspect of the invention there is provided an aqueous disinfecting and cleaning composition in a concentrated form which exhibits reduced irritancy which comprises (preferably consists essentially of):

a disinfecting effective amount of a quaternary ammonium compound having germicidal properties, desirably present in an amount of from about 0.001 - 5% wt.;

10 a mitigating effective amount of at least one nonionic surfactant selected from alkylpolyglycoside compounds;

0.1 - 10%wt. of at least one further nonionic surfactant, preferably selected from alkoxylated primary alcohols;

15 0 - 3%wt. of a polymeric cationic surfactant based on a polyquaternary ammonium salt;

0 - 3%wt. of a builder, desirably present in an amount of about 0.1 - 0.5%wt.;

0 - 5%wt. of one or more conventional additives particularly coloring agents, fragrances and fragrance solubilizers, viscosity modifying agents such as thickeners, pH adjusting agents and pH buffers including organic and inorganic salts; and,

20 water to form 100%wt. of the concentrate form of the inventive compositions.

In accordance with a second aspect of the invention there is provided an aqueous disinfecting and cleaning composition in a concentrated form which exhibits reduced irritancy which comprises (preferably consists essentially of):

25 a disinfecting effective amount of a quaternary ammonium compound having germicidal properties, desirably present in an amount of from about 0.001 - 5% wt.;

a mitigating effective amount of a binary surfactant system which comprises both (a) at least one nonionic surfactant selected from alkylpolyglycoside

30

compounds, with (b) at least one further nonionic surfactant compound which is based on a polymeric alkylene oxide block copolymer, desirably present in an amount of from 0.1 - 10%wt.

0.1 - 10%wt. of at least one further nonionic surfactant, preferably selected from alkoxylated primary alcohols;

0 - 3%wt. of a polymeric cationic surfactant based on a polyquaternary ammonium salt;

0 - 3%wt. of a builder, desirably present in an amount of about 0.1 - 0.5%wt.;

0 - 5%wt. of one or more conventional additives particularly coloring agents, fragrances and fragrance solubilizers, viscosity modifying agents such as thickeners, pH adjusting agents and pH buffers including organic and inorganic salts; and,

water to form 100%wt. of the concentrate form of the inventive compositions.

In accordance with further preferred embodiments of the invention there are provided aqueous dilutions of the concentrated disinfecting and cleaning composition described above, which provides effective cleaning and sanitization.

In certain particularly preferred embodiments, the non-aqueous content of the compositions is particularly low, generally less comprising less than 10%wt., based on the total weight of the composition. Surprisingly however, the compositions provide both effective sanitization and good cleaning.

In a further aspect of the invention there is provided a process for cleaning and/or disinfecting surfaces in need of such treatment which includes contacting a surface with a concentrate composition or aqueous dilution of a concentrate composition as taught herein.

In a still further aspect of the invention there is provided an aqueous hard surface cleaning and sanitizing composition characterized in exhibiting a reduced potential for ocular irritancy which composition contains a quaternary ammonium compound having germicidal properties, said composition further comprising a irritation mitigating effective amount of a nonionic surfactant based on an alkylpolyglycoside.


In a still further aspect of the invention there is provided an aqueous hard surface cleaning and sanitizing composition characterized in exhibiting a reduced potential for ocular irritancy which composition contains a quaternary ammonium compound having germicidal properties, said composition further comprising a irritation  
5 mitigating effective amount of a nonionic surfactant based on an alkylpolyglycoside in combination with a mitigating effective amount of a nonionic surfactant based on a nonionic surfactant compound which is based on a polymeric alkylene oxide block copolymer.

According to a still further aspect of the invention there is provided a process  
10 for mitigating potential ocular irritation caused by an aqueous hard surface cleaning and sanitizing composition which contains a quaternary ammonium compound having germicidal properties, which process includes the step of providing a mitigating effective amount of a nonionic surfactant based on an alkylpolyglycoside to the composition.

According to a yet further aspect of the invention there is provided a process  
15 for mitigating potential ocular irritation caused by an aqueous hard surface cleaning and sanitizing composition which process includes the step of providing a mitigating effective amount of a nonionic surfactant based on an alkylpolyglycoside to an aqueous composition which contains a quaternary ammonium compound having  
20 germicidal properties and at least one nonionic surfactant compound.

In particularly preferred embodiments the concentrated disinfecting and cleaning compositions provided herein provide good cleaning, effective sanitization of surfaces particularly hard surfaces, and low irritancy to the consumer, especially low ocular irritation.


25 The compositions of the invention include a disinfecting effective amount of a quaternary ammonium compound having germicidal properties. Particularly useful quaternary ammonium compounds and salts thereof include quaternary ammonium germicides which may be characterized by the general structural formula:



where at least one of  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  is a hydrophobic, aliphatic, aryl aliphatic or aliphatic aryl radical of from 6 to 26 carbon atoms, and the entire cation portion of the molecule has a molecular weight of at least 165. The hydrophobic radicals may be long-chain alkyl, long-chain alkoxy aryl, long-chain alkyl aryl, halogen-substituted long-chain alkyl aryl, long-chain alkyl phenoxy alkyl, aryl alkyl, etc. The remaining radicals on the nitrogen atoms other than the hydrophobic radicals are substituents of a hydrocarbon structure usually containing a total of no more than 12 carbon atoms. The radicals  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  may be straight chained or may be branched, but are preferably straight chained, and may include one or more amide or ester linkages. The radical X may be any salt-forming anionic radical.

Exemplary quaternary ammonium salts within the above description include the alkyl ammonium halides such as cetyl trimethyl ammonium bromide, alkyl aryl ammonium halides such as octadecyl dimethyl benzyl ammonium bromide, N-alkyl pyridinium halides such as N-cetyl pyridinium bromide, and the like. Other suitable types of quaternary ammonium salts include those in which the molecule contains either amide or ester linkages such as octyl phenoxy ethoxy ethyl dimethyl benzyl ammonium chloride, N-(laurylcocoaminoformylmethyl)-pyridinium chloride, and the like. Other very effective types of quaternary ammonium compounds which are useful as germicides include those in which the hydrophobic radical is characterized by a substituted aromatic nucleus as in the case of lauryloxyphenyltrimethyl ammonium chloride, cetylaminophenyltrimethyl ammonium methosulfate, dodecylphenyltrimethyl ammonium methosulfate, dodecylbenzyltrimethyl ammonium chloride, chlorinated dodecylbenzyltrimethyl ammonium chloride, and the like.

Preferred quaternary ammonium compounds which act as germicides and which are found useful in the practice of the present invention include those which have the structural formula:



wherein R<sub>2</sub> and R<sub>3</sub> are the same or different C<sub>8</sub>-C<sub>12</sub>alkyl, or R<sub>2</sub> is C<sub>12-16</sub>alkyl, C<sub>8</sub>-  
 18alkylethoxy, C<sub>8-18</sub>alkylphenoethoxy and R<sub>3</sub> is benzyl, and X is a halide, for example  
 5 chloride, bromide or iodide, or X may be methosulfate. The alkyl groups recited in R<sub>2</sub>  
 and R<sub>3</sub> may be straight chained or branched, but are preferably substantially linear.

Particularly useful quaternary germicides include compositions which include  
 a single quaternary, as well as mixtures of two or more different quaternaries.


Particularly useful quaternary germicides include BARDAC® 205M, and BARDAC®  
 10 208M or BTC® 885 which is described to be a blend of alkyl dimethyl benzyl  
 ammonium chlorides; BARDAC® 2050 and BARDAC® 2080 or BTC® 818 which  
 is described to be based on dialkyl(C<sub>8</sub>-C<sub>10</sub>)dimethyl ammonium chloride; BARDAC®  
 2250 and BARDAC® 2280 or BTC® 1010 which is described to a composition  
 which includes didecyl dimethyl ammonium chloride; BARDAC® LF and  
 15 BARDAC® LF 80 which is described to be based on dioctyl dimethyl ammonium  
 chloride; BARQUAT® MB-50, HYAMINE® 3500, BARQUAT® MB-80, BTC®  
 835 or BTC 8358 each described to be based on alkyl dimethyl benzyl ammonium  
 chloride; BARQUAT® MX-50, BARQUAT® MX-80, BTC® 824 or BTC® 8248  
 each described to be a composition based on alkyl dimethyl benzyl ammonium  
 20 chloride; BARQUAT® OJ-50, BARQUAT® OJ-80, BTC® 2565, or BTC® 2658  
 each described to be a composition based on alkyl dimethyl benzyl ammonium  
 chloride; BARQUAT® 4250, BARQUAT® 4280, BARQUAT® 4250Z,  
 BARQUAT® 4280Z, BTC® 2125, or BTC® 2125M each described to be a  
 composition based on alkyl dimethyl benzyl ammonium chloride and/or alkyl  
 25 dimethyl ethyl benzyl ammonium chloride; BARQUAT® MS-100 or BTC® 324-P-  
 100 each described to be based on myristyl dimethyl benzyl ammonium chloride;  
 HYAMINE® 2389 described to be based on methyl dodecyl benzyl ammonium  
 chloride and/or methyl dodecyl xylene-bis-trimethyl ammonium chloride;



HYAMINE® 1622 described to be an aqueous solution of benzethonium chloride;  
HYAMINE® 3500-NF or BTC® 50 each described to be based on alkyl dimethyl  
benzyl ammonium chloride; as well as BARQUAT® 1552 or BTC® 776 described  
to be based on alkyl dimethyl benzyl ammonium chloride and/or dialkyl methyl  
benzyl ammonium chloride. (Each of these recited materials are presently  
commercially available from Lonza, Inc., Fairlawn, NJ and/or from Stepan Co.,  
Northfield IL). It is to be understood that these quaternary ammonium compounds  
may be used singly or in mixtures of two or more. These quaternary ammonium  
compounds are desirably present in the concentrate compositions in an amount of  
from about 0.001 - 5% wt., are desirably present in an amount of from 0.1 - 3%wt.  
and most desirably are present in an amount of from 0.5 - 3%wt. When diluted in a  
larger volume of water to form a cleaning and disinfecting composition, the  
quaternary ammonium compounds should be present in sufficient amount such that  
they are in a concentration of at least about 150 parts per million (p.p.m.), more  
desirably at least about 175 p.p.m. and most desirably about 200 - 250 p.p.m. The  
present inventors have surprisingly found that certain of their formulations exhibited  
effective cleaning and disinfecting with less than 200 p.p.m. of the quaternary  
ammonium compounds in cleaning compositions which is an amount below which is  
generally believed to be necessary for disinfecting efficacy.


The compositions of the invention also include a mitigating effective amount  
of at least one nonionic surfactant based on an alkylpolyglycoside compound.  
Exemplary suitable compounds include alkyl monoglycosides and polyglycosides are  
prepared generally by reacting a monosaccharide, or a compound hydrolyzable to a  
monosaccharide with an alcohol such as a fatty alcohol in an acid medium. Various  
glycoside and polyglycoside compounds including alkoxylated glycosides and  
processes for making them are disclosed in U.S. Patent No. 2,974,134; U.S. Patent  
No. 3,219,656; U.S. Patent No. 3,598,865; U.S. Patent No. 3,640,998; U.S. Patent No.  
3,707,535; U.S. Patent No. 3,772,269; U.S. Patent No. 3,839,318; U.S. Patent No.  
3,974,138; U.S. Patent No. 4,223,129; and U.S. Patent No. 4,528,106.

Exemplary alkyl glycoside surfactants suitable for use in the practice of this invention may be represented by formula I below:




wherein:

- 5 R is a monovalent organic radical containing from about 6 to about 30, preferably from about 8 to about 18 carbon atoms;  
 R<sub>1</sub> is a divalent hydrocarbon radical containing from about 2 to about 4 carbon atoms, especially ethyl and propyl radicals;  
 O is an oxygen atom;  
 10 y is a number which has an average value from about 0 to about 1 and is preferably 0;  
 G is a moiety derived from a reducing saccharide containing 5 or 6 carbon atoms; and  
 x is a number having an average value from about 1 to 5 (preferably from 1.1 to 2);  
 15 Z is selected O<sub>2</sub>M<sup>1</sup>,



- O(CH<sub>2</sub>), CO<sub>2</sub>M<sup>1</sup>, OSO<sub>3</sub>M<sup>1</sup>, or O(CH<sub>2</sub>)SO<sub>3</sub>M<sup>1</sup>; R<sub>2</sub> is (CH<sub>2</sub>)CO<sub>2</sub>M<sup>1</sup> or CH=CHCO<sub>2</sub>M<sup>1</sup>; with the proviso that Z can be O<sub>2</sub>M<sup>1</sup> only if Z is in-  
 20 place of a primary hydroxyl group in which the primary hydroxyl-bearing carbon atom,  
 —CH<sub>2</sub>OH, is oxidized to form a




- group);  
 25 b is a number of from 0 to 3x+1 preferably an average of from 0.5 to 2 per glycosal group;  
 p is 1 to 10,

M<sup>1</sup> is H<sup>+</sup> or an organic or inorganic cation, such as, for example, an alkali metal, ammonium, monoethanolamine, or calcium.

As defined in Formula I above, R is generally the residue of a fatty alcohol having from about 8 to 30 and preferably 8 to 18 carbon atoms.

5 Most preferably, the inventive compositions include an alkylpolyglycoside compound according to the structure:



wherein:

10 R is an alkyl group, preferably a linear alkyl chain, which comprises C8 to C16 alkyl groups;

x is an integer value of from 0 – 3, inclusive.

15 Examples of such alkylpolyglycoside compounds according to this structure include: where R is comprised substantially of C8 and C10 alkyl chains yielding an average value of about 9.1 alkyl groups per molecule (Glucopon® 220 UP, Glucopon® 225 DK); where R is comprised of C8, C10, C12, C14 and C16 alkyl chains yielding an average value of about 10.3 alkyl groups per molecule (Glucopon® 425); where R is comprised substantially of C12, C14 and C16 alkyl chains yielding an average value of about 12.8 alkyl groups per molecule (Glucopon® 600 UP, Glucopon® 625 CSUP, and Glucopon® 625 FE, all of which are available from Henkel Corp., Ambler PA.)  
20 Also useful as the alkylpolyglycoside compound is Triton® CG-110 (Union Carbide Corp.).

Further examples of commercially available alkylglycosides as described above include, for example, Glucopon® 325N which is described as being a 50% C<sub>9</sub>-C<sub>11</sub> alkyl polyglycoside, also commonly referred to as D-glucopyranoside (from  
25 Henkel Corp, Ambler PA). Particularly preferred as the alkylpolyglycoside compounds are those illustrated in the Examples.

The inclusion of the alkylpolyglycoside compound to the compositions significantly reduce the irritation potential of the aqueous compositions as compared to like compositions which however omit this constituent. In corresponding copending US Serial No 08/984670 compositions which included only the nonionic surfactant constituent based on a polymeric alkylene oxide block copolymer were found to have generally satisfactory low ocular irritation potential. Therein, alkylpolyglycosides are indicated as one of a number of optional nonionic surfactants which might also be present, but that application does not indicate that any of these nonionic surfactants might be useful in providing an irritation mitigating effect. Only presently have the present inventors discovered that compositions, especially those according to particularly preferred embodiments of the present invention, which further include the alkylpolyglycoside compound as a necessary constituent have even further reduced ocular irritation potential. While not wishing to be bound by the following, it is theorized that the presence of both a nonionic surfactant constituent based on a compound other than an alkylpolyglycoside in conjunction with the nonionic surfactant based on the alkylpolyglycoside compound have a synergistic or complementary effect in reducing the irritation potential of such aqueous compositions. When both are present, as in the first embodiment of the invention, the former to the latter are desirably included in relative weight ratios of from 3:1 to 1.5:1. This irritation mitigating effect has also been found in the inventive compositions according to the second aspect of the invention, wherein there is present the binary surfactant system which comprises at least one nonionic surfactant selected from alkylpolyglycoside compounds, and at least one further nonionic surfactant compound which is based on a polymeric alkylene oxide block copolymer. According to this second embodiment of the invention, generally good results have been observed when the weight ratio of the alkylpolyglycoside compounds to the polymeric alkylene oxide block copolymer are in respective ratios of from 0.5:1 – 2:1.

The amounts of the nonionic surfactant based on an alkylpolyglycoside compound to the compositions may vary in accordance with the level of irritancy mitigation sought. Generally, the alkylpolyglycoside surfactant may be included in

any amount which mitigates irritancy. Good efficacy is found when present from about 0.01 – 10%wt. based on the total weight of the composition, but amounts of from 0.1 – 7%wt. and preferably from about 0.5 – 4%wt. are preferred. It is to be understood that the amount which is to be included will vary upon several factors such as the amounts of the other constituents present in a composition, as well as the irritancy levels of such other constituents. The optimal amount of the alkylpolyglycoside surfactant to be included may be determined by routine experimentation, such as by the method outlined with reference to the Examples.

In compositions according to the second aspect of the invention, the amounts of the two surfactants which comprise the binary surfactant system, may vary in accordance with the level of irritancy mitigation sought. Generally, the at least one nonionic surfactant selected from alkylpolyglycoside compounds is present from about 0.01 – 9.99%wt., but is preferably present from 0.01 – 4%wt., and at the same time the nonionic surfactant compound based on the polymeric alkylene oxide block copolymer may also be present in amounts of from 0.01 – 9.99%wt, but preferably are present in an amount of from 0.5 – 2%wt. The optimal amount of the two surfactants which make up the binary surfactant system which are to be included in the inventive compositions may be determined by routine experimentation, such as by the method outlined with reference to the Examples. While not wishing to be bound by the following, it is believed that the presence of the alkylpolyglycoside compounds with the polymeric alkylene oxide block copolymer compounds provide a synergistic improvement with respect to the mitigation of the level of irritancy, particularly the ocular irritancy of the inventive compositions.

By way of non-limiting example, useful nonionic surfactants based on polymeric alkylene oxide block copolymers which are included in the second aspect of the invention, include nonionic surfactants in which the major portion of the molecule is made up of block polymeric C<sub>2</sub>-C<sub>4</sub> alkylene oxides. Such nonionic surfactants, while preferably built up from an alkylene oxide chain starting group, and can have as a starting nucleus almost any active hydrogen containing group including, without limitation, amides, phenols, thiols and secondary alcohols.

One group of such useful nonionic surfactants containing the characteristic alkylene oxide blocks are those which may be generally represented by the formula (A):



where EO represents ethylene oxide,

PO represents propylene oxide,

y equals at least 15,

(EO)<sub>x+z</sub> equals 20 to 50% of the total weight of said compounds, and,

the total molecular weight is preferably in the range of about 2000 to 15,000.

Another group of nonionic surfactants appropriate for use in the new compositions can be represented by the formula (B):



wherein R is an alkyl, aryl or aralkyl group, where the R group contains 1 to 20 carbon atoms, the weight percent of EO is within the range of 0 to 45% in one of the blocks a, b, and within the range of 60 to 100% in the other of the blocks a, b, and the total number of moles of combined EO and PO is in the range of 6 to 125 moles, with 1 to 50 moles in the PO rich block and 5 to 100 moles in the EO rich block.

Further nonionic surfactants which in general are encompassed by Formula B include butoxy derivatives of propylene oxide/ethylene oxide block polymers having molecular weights within the range of about 2000-5000.

Still further useful nonionic surfactants containing polymeric butoxy (BO) groups can be represented by formula (C) as follows:



wherein R is an alkyl group containing 1 to 20 carbon atoms,


n is about 5-15 and x is about 5-15.

Also useful as the nonionic block copolymer surfactants, which also include polymeric butoxy groups, are those which may be represented by the following formula (D):



wherein n is about 5-15, preferably about 15,  
x is about 5-15, preferably about 15, and  
y is about 5-15, preferably about 15.

Still further useful nonionic block copolymer surfactants include ethoxylated derivatives of propoxylated ethylene diamine, which may be represented by the following formula:



where (EO) represents ethoxy,  
(PO) represents propoxy,

the amount of (PO)<sub>x</sub> is such as to provide a molecular weight prior to ethoxylation of about 300 to 7500, and the amount of (EO)<sub>y</sub> is such as to provide about 20% to 90% of the total weight of said compound.

Of these, the most preferred are those which are represented by formula (A) above; specific examples of which include those materials presently commercially available under the tradename "Pluronic®", and in particular the Pluronic® F series, Pluronic® L series, Pluronic® P series, as well as in the Pluronic® R series, each of which are generally described to be block copolymers of propylene oxide and ethylene oxide. Generally those of the Pluronic® L series and the Pluronic® R series are preferred as these are supplied in liquid form by the manufacturer and are readily formulated into the present inventive compositions. These are also available in a wide range of HLB values, and those having HLB values in the range of 1.0 - 23.0 may be used, although those with intermediate HLB values such as from about 12.0 - 18.0 are found to be particularly advantageous. These materials are presently commercially available from BASF AG (Ludwigshafen, Germany) as well as from BASF Corp. (Mt. Olive Township, New Jersey).

Other useful exemplary nonionic block copolymers based on a polymeric ethoxy/propoxy units which may also be used include those presently commercially available in the Poly-Tergent® E, and Poly-Tergent® P series of materials from Olin

Chemicals Corp., (Stamford CT). These are described to be nonionic surfactants based on ethoxy/propoxy block copolymers, conveniently available in a liquid form from its supplier.

The inventive compositions further include at least one further nonionic surfactant which provides a further deterative benefit to the inventive compositions. Preferred nonionic surfactants provide surprisingly good levels of cleaning performance, particularly in conjunction with the preferred quaternary ammonium compounds described herein.

A preferred class of nonionic surfactants are alkoxyated alcohols. These include the condensation products of a higher alcohol (e.g., an alkanol containing about 8 to 18 carbon atoms in a straight or branched chain configuration) condensed with about 2 to 30 moles of ethylene oxide, for example, lauryl or myristyl alcohol condensed with about 16 moles of ethylene oxide, tridecanol condensed with about 6 to moles of ethylene oxide, myristyl alcohol condensed with about 10 moles of ethylene oxide per mole of myristyl alcohol, the condensation product of ethylene oxide with a distillation or separation fraction of coconut fatty alcohol containing a mixture of fatty alcohols with alkyl chains varying from 10 to about 14 carbon atoms in length and wherein the condensate contains either about 6 moles of ethylene oxide per mole of total alcohol or about 9 moles of ethylene oxide per mole of alcohol and tallow alcohol ethoxylates containing 6 ethylene oxide to 11 ethylene oxide per mole of alcohol.

A preferred group of the foregoing nonionic surfactants are the Neodol® ethoxylates (Shell Chemical Co., Houston TX); which are higher aliphatic, primary alcohols containing about 9-15 carbon atoms, such as a C<sub>11</sub> alkanol condensed with 7 moles of ethylene oxide (Neodol® 1-7), C<sub>9</sub>-C<sub>11</sub> alkanol condensed with an average of 2.5 moles of ethylene oxide (Neodol® 91-2.5); C<sub>9</sub>-C<sub>11</sub> alkanol condensed with 6 moles of ethylene oxide (Neodol® 91-6), C<sub>9</sub>-C<sub>11</sub> alkanol condensed with 8 moles of ethylene oxide (Neodol® 91-8), C<sub>12-13</sub> alkanol condensed with 6.5 moles ethylene oxide (Neodol® 23-6.5), C<sub>12-13</sub> alkanol condensed with 7 moles ethylene oxide (Neodol® 23-7), C<sub>12-15</sub> alkanol condensed with 7 moles of ethylene oxide (Neodol®



25-7),  $C_{12-15}$  alkanol condensed with 9 moles ethylene oxide (Neodol® 25-9),  $C_{12-15}$  alkanol condensed with 12 moles ethylene oxide (Neodol® 25-12),  $C_{14-15}$  alkanol condensed with 13 moles ethylene oxide (Neodol® 45-13), and the like.


5 A further preferred group of nonionic surfactants which are advantageously present in the inventive compositions are those presently marketed under the Genapol® tradename. Particularly useful are those in the Genapol® "26-L" series which include for example: C12-16 linear alcohols condensed with 1 mole of ethylene oxide (Genapol® 24-L-3); C12-16 linear alcohols condensed with 1.6 moles of ethylene oxide (Genapol® 26-L-1.6); C12-16 linear alcohols condensed with 2 moles of ethylene oxide (Genapol® 26-L-2); C12-16 linear alcohols condensed with 3 moles of ethylene oxide (Genapol® 26-L-3); C12-16 linear alcohols condensed with 5 moles of ethylene oxide (Genapol® 26-L-5); as well as C12-16 linear alcohols condensed with varying amounts of ethylene oxide to provide specific cloud points of the surfactant (i.e., Genapol® 26-L-60, Genapol® 26-L-60N, and Genapol® 26-L-98N).  
10 These materials are commercially available from a variety of sources, including Clariant Corp. (Charlotte, N.C.).

Additional useful nonionic surfactants include those based on alcohol and ethylene oxide condensates of a secondary aliphatic alcohol. These alcohols contain 8 to 18 carbon atoms in a straight or branched chain configuration and are condensed with 5 to 30 moles of an alkylene oxide, especially ethylene oxide. Examples of  
20 commercially available nonionic detergents of the foregoing type are  $C_{11}-C_{15}$  secondary alkanols condensed with either 9 ethylene oxides (Tergitol® 15-S-9) or 7 ethylene oxides (Tergitol® 15-S-7) marketed by Union Carbide Corp. (Danbury CT). It is to be understood that these nonionic alkoxylated secondary alcohol surfactant compounds may be used singly or in mixtures of two or more such compounds.


25 Further useful nonionic surfactants include certain alkoxylated linear aliphatic alcohol surfactants which are believed to be the condensation products of a  $C_8-C_{10}$  hydrophilic moiety with alkylene oxides, especially polyethylene oxide and or polypropylene oxide moieties. Such alkoxylated linear alcohol surfactants are  
30 presently commercially available under the tradename PolyTergent® (Olin Chemical

Co., Stamford CT). Of these particularly useful are those which are marketed as PolyTergent® SL-22, PolyTergent® SL-42, PolyTergent® SL-62 and PolyTergent® SL-29, of which PolyTergent® SL-62 is particularly advantageous. PolyTergent® SL-92 is described as being a moderately foaming, biodegradable alkoxyated linear alcohol surfactant having on average 8 moles of oxyethylene groups per molecule. These alkoxyated linear alcohol surfactants provide good deterative action in the removal of many types of fats and greases such as are frequently found in soils on hard surfaces, as well as providing a further solubilizing effects and may be included in the concentrate compositions according to the present invention with advantage. The preferred alkoxyated linear alcohol surfactants also exhibit low levels of ocular irritation in the concentrate compositions.

Further useful nonionic surfactants include alkoxyated, and particularly ethoxyated octyl and nonyl phenols according to the following general structural formulas:



or,



in which the  $\text{C}_9\text{H}_{19}$  group in the latter formula is a mixture of branched chained isomers. In both formulae, x indicates an average number of ethoxy units in the side chain. Suitable non-ionic ethoxyated octyl and nonyl phenols include those having from about 7 to about 13 ethoxy units. Such compounds are commercially available under the trade name Triton® X (Union Carbide, Danbury CT).

Exemplary alkoxyated alkyl phenols useful as a nonionic surfactant also include certain compositions presently commercially available from the Rhône-Poulenc Co., (Cranbury, NJ) under the general trade name Igepal®, which are described to be octyl and nonyl phenols. These specifically include Igepal® CO730

which is described as an ethoxylated nonyl phenol having an average of 15 ethoxy groups per molecule.

5 These nonionic surfactant compounds described above may be used singly or in mixtures. When present, these further nonionic surfactants comprise 0.01 - 10%wt. of the concentrate compositions, desirably comprise 0.1 - 8%wt. and most desirably  
comprise about 2 - 6%wt. and especially about 5%wt. of the concentrate compositions taught herein.

10 The inventive compositions optionally but desirably include a builder. Such a builder constituent may be present in an amount of from 0 - 3%wt. but preferably 0.1 - 0.5%wt. based on the total weight of the concentrate compositions taught herein. Such include water soluble inorganic builders which can be used alone, in admixture with other water soluble inorganic builders, as well as in conjunction with one or more organic alkaline sequestrant builder salts.


15 Exemplary builders include alkali metal carbonates, phosphates, polyphosphates and silicates. More specific examples include sodium tripolyphosphate, sodium carbonate, sodium bicarbonate, sodium tetraborate, potassium carbonate, sodium polyphosphate, potassium pyrophosphate, potassium tripolyphosphate, and sodium hexametaphosphate. Further exemplary builders also include organic alkaline sequestrant builder salts such as alkali metal polycarboxylates  
20 including water-soluble citrates such as sodium and potassium citrate, sodium and potassium tartarate, sodium and potassium ethylenediaminetetraacetate, sodium and potassium N-(2-hydroxyethyl)-ethylene diamine triacetates, sodium and potassium nitrilotriacetates, as well as sodium and potassium tartrate mono- and di-succinates. Also useful are gluconate or glucoheptonate salts particularly sodium gluconate and  
25 sodium glucoheptonate. Particularly advantageously used are di-, tri- and tetrasodium salts of ethylenediaminetetraacetic acid, especially tetrasodium salts thereof. As noted, these organic builder salts may be used individually, as a combination of two or more organic builder salts, as well as in conjunction with one or more detergency builders, including those indicated above. It is to be understood that many of these

builder materials also provide a useful pH stabilizing effect in the compositions in which they form a part.

As is noted above, the compositions according to the invention are aqueous in nature. Water is added to the constituents in order to provide 100% by weight of the composition. The water may be tap water, but is preferably distilled and is most preferably deionized water. If the water is tap water, it is preferably substantially free of any undesirable impurities such as organics or inorganics, especially mineral salts which are present in hard water which may thus interfere with the operation of one or more of the constituents of the aqueous compositions according to the invention.

Preferably the concentrate composition comprise at least 80% water.

An optional but in certain instances, particularly desirable further constituent is a cationic polymeric polyquaternary ammonium salt, especially a halogen salt such as a chloride salt. Such a material is one which includes at least one repeating monomer unit wherein such monomer includes as part of its structure a quaternary ammonium. A particularly useful class of such materials are those sold under the trade designation "Mirapol®" and are available from Rhône-Poulenc Surfactant & Specialty Chemicals Co. (Cranbury, NJ). These materials are highly cationic in nature, and are believed to be in accordance with the following general structure:



wherein n is an integer or 2 or greater, and is desirably in the range of 2 - 12, more desirably is about 6. Such a material is commercially available as Mirapol® A-15 from Rhône-Poulenc, identified above. When present, this material may comprise to 3%wt.

The inventors have found that the inclusion of such material provides a useful soil suspending benefit which is desirable from a cleaning standpoint, although it has also been observed by the inventors that inclusion of such a material may have a detrimental effect on the disinfecting properties of the compositions.

The constituents which may be used in the compositions according to the invention are known, many of which are described in *McCutcheon's Detergents and Emulsifiers*, North American Edition, 1991; *Kirk-Othmer, Encyclopedia of Chemical Technology*, 3rd Ed., Vol. 22, pp. 346-387, the contents of which are herein  
5 incorporated by reference.

The compositions according to the invention are useful in the disinfecting and/or cleaning of surfaces, especially hard surfaces in need of such treatment. These in particular include surfaces wherein the presence of gram positive and/or gram negative bacteria are suspected. In accordance with the present inventive process,  
10 cleaning and/or disinfecting of such surfaces comprises the step of applying a stain releasing and a disinfecting effective amount of a composition as taught herein to such a stained surface. Afterwards, the compositions are optionally but desirably wiped, scrubbed or otherwise physically contacted with the hard surface, and further optionally, may be subsequently rinsed from such a cleaned and disinfected hard  
15 surface.

Such a hard surface cleaning and disinfecting composition according to the invention is may be provided as a ready to use product which may be directly applied to a hard surface, but is desirably provided in a concentrated form intended to be diluted in water to form a cleaning composition therefrom.

By way of example, hard surfaces include surfaces composed of refractory materials such as: glazed and unglazed tile, porcelain, ceramics as well as stone including marble, granite, and other stones surfaces; glass; metals; plastics e.g. polyester, vinyl; fiberglass, Formica®, Corian® and other hard surfaces known to the industry. Hard surfaces which are to be particularly denoted are lavatory fixtures such  
20 as shower stalls, bathtubs and bathing appliances (racks, shower doors, shower bars) toilets, bidets, wall and flooring surfaces especially those which include refractory materials and the like. Further hard surfaces which are to be denoted are those associated with kitchen environments and other environments associated with food  
25 preparation, including cabinets and countertop surfaces as well as walls and floor

surfaces especially those which include refractory materials, plastics, Formica®, Corian® and stone.

The hard surface cleaner composition provided according to the invention can be also be provided as a ready to use product in a manually operated spray dispensing container. Such a typical container is generally made of synthetic polymer plastic material such as polyethylene, polypropylene, polyvinyl chloride or the like and includes spray nozzle, a dip tube and associated pump dispensing parts and is thus ideally suited for use in a consumer "spray and wipe" application. In such an application, the consumer generally applies an effective amount of the cleaning composition using the pump and within a few moments thereafter, wipes off the treated area with a rag, towel, or sponge, usually a disposable paper towel or sponge. In certain applications, however, especially where undesirable stain deposits are heavy, the cleaning composition according to the invention may be left on the stained area until it has effectively loosened the stain deposits after which it may then be wiped off, rinsed off, or otherwise removed. For particularly heavy deposits of such undesired stains, multiple applications may also be used.

In a yet a further embodiment, the compositions according to the invention may be formulated so that it may be useful in conjunction with a "aerosol" type product wherein it is discharged from a pressurized aerosol container. If the inventive compositions are used in an aerosol type product, it is preferred that corrosion resistant aerosol containers such as coated or lined aerosol containers be used. Such are preferred as they are known to be resistant to the effects of basic formulations. Known art propellants such as liquid propellants as well as propellants of the non-liquid form, i.e., pressurized gases, including carbon dioxide, air, nitrogen, hydrocarbons as well as others may be used. Also, while satisfactory for use, fluorocarbons may be used as a propellant but for environmental and regulatory reasons their use is preferably avoided. In such an embodiment, the cleaning composition is dispensed by activating the release nozzle of said aerosol type container onto the stain and/or stain area, and in accordance with a manner as above-described a stain is treated and removed.

Whereas the present invention is intended to be used in the types of liquid forms described, the compositions according to the invention are desirably diluted with a further amount of water to form a cleaning and disinfecting solution therefrom. In such a proposed diluted cleaning solution, the greater the proportion of water added to form said cleaning dilution, the greater may be the reduction of the rate and/or efficacy of the thus formed cleaning solution in the cleaning of a hard surface, as well as a reduction in disinfectant efficacy. Accordingly, longer residence times upon the stain to effect their loosening and/or the usage of greater amounts may be necessitated. Conversely, nothing in the specification shall be also understood to limit the forming of a "super-concentrated" cleaning composition based upon the composition described above. Such a super-concentrated composition is essentially the same as the compositions described above except in that they include a lesser amount of water.

While the cleaning compositions are most beneficial for use in their form, i.e., their form as described above, they may also be diluted to form a cleaning composition therefrom. Such cleaning compositions may be easily prepared by diluting measured amounts of the compositions in further amounts of water by the consumer or other end user in certain weight ratios of composition: water, and optionally, agitating the same to ensure even distribution of the composition in the water. The concentrate compositions according to the invention may be used without further dilution, but may also be used with a further aqueous dilution, i.e., in concentrate composition: water concentrations of 1:0, to extremely dilute dilutions such as 1:1000. When subjected to further aqueous dilution, such a dilution is preferably a weight or volume ratio proportion of from 1:10 - 1:64, and most desirably is about 1:64. The actual dilution selected is in part determinable by the degree and amount of dirt and grime to be removed from a surface(s), the amount of mechanical force imparted to remove the same, as well as the observed efficacy of a particular dilution. Generally better results and faster removal is to be expected at lower relative dilutions of the composition and the water.

Other conventional optional additives, although not particularly elucidated herein may also be included in the present inventive compositions. Exemplary optional conventional additives include but are not limited to: pH adjusting agents and pH buffers including organic and inorganic salts; non-aqueous solvents, perfumes, perfume carriers, optical brighteners, coloring agents such as dyes and pigments, opacifying agents, hydrotropes, antifoaming agents, viscosity modifying agents such as thickeners, enzymes, anti-spotting agents, anti-oxidants, anti-corrosion agents as well as others not specifically elucidated here. These should be present in minor amounts, preferably in total comprise less than about 5% by weight of the compositions, and desirably less than about 3%wt.

The following examples below illustrate exemplary and preferred formulations of the concentrate composition according to the instant invention. It is to be understood that these examples are presented by means of illustration only and that further useful formulations fall within the scope of this invention and the claims may be readily produced by one skilled in the art and not deviate from the scope and spirit of the invention.

Throughout this specification and in the accompanying claims, weight percents of any constituent are to be understood as the weight percent of the active portion of the referenced constituent, unless otherwise indicated.

#### Example Formulations:

##### Preparation of Example Formulations:

Exemplary formulations illustrating certain preferred embodiments of the inventive compositions and described in more detail in Table 1 below were formulated generally in accordance with the following protocol.

Into a suitably sized vessel, a measured amount of water was provided after which the constituents were added in no specific or uniform sequence, which indicated that the order of addition of the constituents was not critical. All of the constituents were supplied at room temperature, and any remaining amount of water was added thereafter. Certain of the nonionic surfactants if gels at room temperature



were first preheated to render them pourable liquids prior to addition and mixing. Mixing of the constituents was achieved by the use of a mechanical stirrer with a small diameter propeller at the end of its rotating shaft. Mixing, which generally lasted from 5 minutes to 120 minutes was maintained until the particular exemplary formulation appeared to be homogeneous. The exemplary compositions were readily pourable, and retained well mixed characteristics (i.e., stable mixtures) upon standing for extend periods. The compositions of the example formulations are listed on Table 1. The weight percentages indicated the "as supplied" weights of the named constituent.

As is indicated, to all of the formulations of Table 1 was added sufficient deionized water in "quantum sufficient" to provide 100 parts by weight of a particular formulation.

Table 1	Control	542-050D
	Comp.1	Ex.1
Neodol® 25-7	5.00	—
PolyTergent® SL-62	—	4.00
BTC® 8358 (80%)	1.625	1.625
Pluronic L64	2.0	—
Glucopon® 325N (50%)	—	4.0
EDTA (38%)	0.25	0.25
Fragrance	0.20	0.20
Dye Solution	0.20	0.20
DI water	q.s.	q.s.

5 The identity of the constituents of Table 1 above are described in more detail on Table 2, below, including the "actives" percentage of each were a constituent was not 100%wt. "actives".

TABLE 2	
constituent:	identity:
Pluronic® L-64	nonionic ethoxy/propoxy block copolymer surfactant (BASF Corp.)
Neodol® 25-7	nonionic C12-15 alkanol condensed with 7 moles ethylene oxide (Shell Chemical Co.)
Polytergent ® SL-62	alkoxylated alcohol (BASF Inc.)
Glucopon® 325N (50%)	technical grade mixture of C9-11 alkylpolyglycosides (Henkel Corp.)
BTC® 8358 (80%)	alkyl dimethyl benzyl ammonium chloride (Stepan Co.)
EDTA (38%)	tetrasodium ethylenediaminetetraacetate
Fragrance	proprietary composition
dye solution	proprietary composition
DI water	deionized water

#### Evaluation of Antimicrobial Efficacy:

10 Several of the exemplary formulations described in more detail on Table 1 above were evaluated in order to evaluate their antimicrobial efficacy against

*Staphylococcus aureus* (gram positive type pathogenic bacteria) (ATCC 6538), and  
*Salmonella choleraesuis* (gram negative type pathogenic bacteria) (ATCC 10708)  
The testing was performed in accordance with the protocols outlined in "Use-Dilution  
Method", Protocols 955.14, 955.15 and 964.02 described in Chapter 6 of "Official  
5 Methods of Analysis", 16<sup>th</sup> Edition, of the Association of Official Analytical  
Chemists; "Germicidal and Detergent Sanitizing Action of Disinfectants", 960.09  
described in Chapter 6 of "Official Methods of Analysis", 15<sup>th</sup> Edition, of the  
Association of Official Analytical Chemists; or American Society for Testing and  
Materials (ASTM) E 1054-91 the contents of which are herein incorporated by  
10 reference. This test is also commonly referred to as the "AOAC Use-Dilution Test  
Method".

As is appreciated by the skilled practitioner in the art, the results of the AOAC  
Use-Dilution Test Method indicates the number of test substrates wherein the tested  
organism remains viable after contact for 10 minutes with at test disinfecting  
15 composition / total number of tested substrates (cylinders) evaluated in accordance  
with the AOAC Use-Dilution Test. Thus, a result of "0/60" indicates that of 60 test  
substrates bearing the test organism and contacted for 10 minutes in a test disinfecting  
composition, 0 test substrates had viable (live) test organisms at the conclusion of the  
test. Such a result is excellent, illustrating the excellent disinfecting efficacy of the  
20 tested composition. Results for lesser amount of test substrates such as for 10, 20, 30  
or 40 test substrates provide useful screening results, although insufficient to satisfy  
the requirement of 60 test substrates as dictated by the AOAC Use-Dilution Test.

Results of the antimicrobial testing are indicated on Table 3, below. The  
reported results indicate the number of test cylinders with live test organisms/number  
25 of test cylinders tested for each example formulation and organism tested.

Table 3		
Formulation:	<i>Staphylococcus aureus</i>	<i>Salmonella choleraesuis</i>
Comp.1	0/30	0/30
Ex.1	0/60	1/60

Evaluation of Ocular Irritation:

The ocular irritation characteristics of formulations according to the invention were evaluated using the known Draize Eye test protocol. Evaluation was performed on several formulations according to the invention and described more fully in Table 1 above.

As known to those skilled in the art, the Draize Eye Test measures eye irritation for the grading of severity of ocular lesions, measuring three dimensions: scores obtained for the cornea, iris and conjunctiva. For the cornea, after exposure to the composition, A the cornea opacity is graded on a scale from 1 to 4; B the area of cornea involved is graded on a scale from 1-4 (where the score =  $A \times B \times 5$  may be a total maximum of 80). For evaluation of the iris, after exposure the composition, A the involvement of the iris is graded on a scale of 1-2 (where the score =  $A \times 5$  may be a total maximum of 10). For a evaluation of the conjunctive, A Redness is graded on a scale of 1-3; B Chemosis is graded on a scale of 1-4; and C Discharge is measured on a scale of 1-3 [where the score =  $(A + B + C) \times 2$  may be a maximum of 20]. The maximum total score is the sum of all scores obtained for the cornea, iris and conjunctive (a maximum of 110).

The results of the Draize test are reported below. These indicate that an EPA classification Category "3" was appropriate, where corneal involvement or irritation cleared in "21" days or less. These results are in accordance with the guidelines of the Environmental Protection Agency (EPA), 40 C.F.R. Ch.1, §162.10, (1986).

TABLE 4	
Formulation:	Corneal opacity in test subjects / number of days
Control 1	8.33 / 21
Ex.1	0 / 21

As may be seen from these results, the composition according to Ex. 1 exhibited very low levels of ocular irritation, as is demonstrated by the rapid rate at which corneal opacity cleared in the test subjects.

#### Evaluation of Cleaning Efficacy:

Various formulations amongst those listed above were evaluated for their cleaning efficacy on tile surfaces utilizing the following protocols. "Standard soiled tiles" were prepared for use in the tests. These were prepared in accordance with the protocol described in ASTM 4488-87, Annex A5 "Particulate and Oily Soil/Vinyl Tiles Test Method". This preparation of standard soiled tiles and cleaning protocol was performed for certain of cleaning compositions formed from the formulations described in more detail on Table 1.

Evaluation was performed utilizing a Gardner Washability Apparatus, using a standard soil tiles prepared in accordance with the protocol described above at a standard pressure and sponge stroke settings in order to determine or quantify the cleaning efficiency of the formulations. These formulations were used formed into a cleaning composition wherein 1 part of a formulation of Table 1 was diluted with 64 parts water. For comparative purposes, a 1:64 dilution of a commercially available concentrated cleaning and disinfecting preparation, Lysol® Deodorizing Cleaner "Country Scent" variety was also prepared and evaluated in the same test. In determining the cleaning efficiency of each of the formulations, reflectance values were determined using a Minolta Chromameter where each tile was measured three times and the mean reflectance value are reported below on Table 5. For each of these tiles, there were at least four replicates, each of which were evaluated and used to determine the mean reflectance value of Table 5. Testing was performed for each of the following: a clean unsoiled tile, a soiled tile, and a soiled tile following Gardner Washability Apparatus scrubbing. Such reflectance values were then employed to calculate cleaning efficiency according to the following formula:

$$\text{Cleaning Efficiency} = \frac{L_t - L_s}{L_o - L_s}$$

wherein:

$L_t$  = reflectance average after scrubbing solid tile;

$L_s$  = reflectance average before cleaning soiled tile;

$L_o$  = reflectance average original tile before soiling.

The evaluation procedure noted above was performed in groups of test tiles, wherein the cleaning compositions formed from formulations according to Table 1 were compared to a tiles treated with the cleaning composition formed using a commercially available product, Lysol® Deodorizing Cleaner "Country Scent" formulation (Reckitt & Colman Inc., Montvale NJ). These cleaning efficiency results are shown in the Table 5, following.

Table 5	
Formulation:	Oily
Lysol® Deodorizing Cleaner "Country Scent" formulation	45 - 50%
Comp.1	35 - 40%
Ex.1	50 - 55%

As shown, the measurement of the cleaning effectiveness of the test samples involved the ability of the cleaning composition to remove the test soil from the test substrate. This was expressed by Cleaning Efficiency; as numerical values for a Cleaning Efficiency increase, higher cleaning effectiveness is achieved for the cleaning composition tested. As the results show, the inventive composition showed excellent cleaning characteristics which favorably compare to the commercially available products.

Claims:

- 5        1.     Aqueous disinfecting and cleaning composition which exhibits reduced irritancy  
which comprises,  
a disinfecting effective amount of a quaternary ammonium compound having  
germicidal properties;  
a mitigating effective amount of at least one nonionic surfactant selected from  
10       alkylpolyglycoside compounds;  
0.1 - 10%wt. of a nonionic surfactant;  
0 - 3%wt. of a polymeric cationic surfactant based on a polyquaternary  
ammonium salt;  
0 - 3%wt. of a builder 0 - 5%wt. of one or more conventional additives  
15       particularly coloring agents, fragrances and fragrance solubilizers, viscosity  
modifying agents such as thickeners, pH adjusting agents and pH buffers  
including organic and inorganic salts; and, water to form 100%wt.
- 20       2.     The concentrate composition according to claim 1 wherein the quaternary  
ammonium compound having germicidal properties is present in an amount of  
from about 0.001 - 5% wt.
- 25       3.     The concentrate composition according to claim 1 wherein the at least one  
further nonionic surfactant is present in an amount of from about at 0.1 -  
8%wt.
- 30       4.     The concentrate composition according to claim 3 wherein the at least one  
further nonionic surfactant is an alkoxylated primary alcohol.

5. The concentrate composition according to claim 3 wherein the at least one further nonionic surfactant is a polymeric alkylene oxide block copolymer.
6. An aqueous composition which comprises 1 part of the aqueous disinfecting and cleaning concentrate composition per 10 to 64 parts water.
7. Aqueous disinfecting and cleaning composition in a concentrated form which exhibits reduced irritancy which comprises:  
a disinfecting effective amount of a quaternary ammonium compound having germicidal properties;  
a mitigating effective amount of a binary surfactant system which comprises both (a) at least one nonionic surfactant selected from alkylpolyglycoside compounds, with (b) at least one further nonionic surfactant compound which is based on a polymeric alkylene oxide block copolymer;  
0.1 - 10%wt. of at least one further nonionic surfactant;  
0 - 3%wt. of a polymeric cationic surfactant based on a polyquaternary ammonium salt;  
0 - 3%wt. of a builder;  
0 - 5%wt. of one or more conventional additives particularly coloring agents, fragrances and fragrance solubilizers, viscosity modifying agents such as thickeners, pH adjusting agents and pH buffers including organic and inorganic salts; and,  
water to form 100%wt. of the concentrate form of the inventive compositions.
8. The concentrate composition according to claim 7 wherein the quaternary ammonium compound having germicidal properties is present in an amount of from about 0.001 - 5% wt.
9. The concentrate composition according to claim 7 wherein the binary surfactant system is present in an amount of from 0.1 - 10%wt.



10. The concentrate composition according to claim 7 which comprises at least one further nonionic surfactant.
- 5 11. The concentrate composition according to claim 10 wherein the at least one further nonionic surfactant is an alkoxylated primary alcohol.
- 10 10. An aqueous composition which comprises 1 part of the aqueous disinfecting and cleaning concentrate composition according to claim 7 per 10 to 64 parts water.
11. A process for cleaning and/or disinfecting of hard surfaces which comprises the step of:  
applying an effective amount of a composition according to claim 1 to  
15 the surface.
12. A process for cleaning and/or disinfecting of hard surfaces which comprises the step of:  
applying an effective amount of a composition according to claim 7 to  
20 the surface.
13. A concentrate composition according to claim 1 substantially as described with reference to one or more of the Examples.



The  
Patent  
Office

38

Application No: GB 9807664.9  
Claims searched: 1 to 13

Examiner: Michael Conlon  
Date of search: 16 July 1998

**Patents Act 1977**  
**Search Report under Section 17**

**Databases searched:**

UK Patent Office collections, including GB, EP, WO & US patent specifications, in:  
UK CI (Ed.P): C5D (DHX, DHZ, DJX, D162)  
Int CI (Ed.6): C11D 3/00  
Other: Online: WPI, CLAIMS

**Documents considered to be relevant:**

Category	Identity of document and relevant passage	Relevant to claims
X	GB2309706 A (Reckitt & Colman) Examples 1, 4 and 5	1 at least
X	WO95/31962 A1 (Gojo) Table XIV on page 38	1 at least
X	US4755327 (Bernaducci) Table I and column 6 line 13	1 at least

X	Document indicating lack of novelty or inventive step	A	Document indicating technological background and/or state of the art.
Y	Document indicating lack of inventive step if combined with one or more other documents of same category.	P	Document published on or after the declared priority date but before the filing date of this invention.
&	Member of the same patent family	E	Patent document published on or after, but with priority date earlier than, the filing date of this application.